


1. PET (polyethylene terephthalate) has a higher elastic modulus value than nylon and polyethylene. Engineering analysis D00295173. Data on file.


Clinical strength

Hercules balloons have the dilating strength to relieve your patient's stricture in the GI tract, including the esophagus, pylorus, duodenum, and colon, and may support longer-term luminal patency.



Rapid deflation²

Rapid deflation is achieved by simply applying negative pressure. Even if removal begins prior to complete deflation, the Hercules balloon can still be withdrawn into the endoscope.



Hercules[®] 3 Stage Balloon

Used to endoscopically dilate strictures of the esophagus.

Order Number	Reference Part Number	Catheter Fr	Catheter Length cm	Balloon Diameter mm	Balloon Fr	Balloon Length cm	Pressure atm	Pressure psi	Minimum Accessory Channel Diameter mm
G31928	HBD-8-9-10	7	180	8-9-10	24-27-30	8	4-6-8	60-90-120	2.8
G31925	HBD-10-11-12	7	180	10-11-12	30-33-36	8	2-4-6	30-60-90	2.8
G31926	HBD-12-13.5-15	7	180	12-13.5-15	36-40.5-45	8	2-4-6	30-60-90	2.8
G48732	HBD-15-16.5-18	7	180	15-16.5-18	45-49.5-54	8	2-4-6	30-60-90	2.8
G31927	HBD-18-19-20	7	180	18-19-20	54-57-60	8	2-4-6	30-60-90	2.8

Hercules[®] 3 Stage Wire Guided Balloon

Used to endoscopically dilate strictures of the gastrointestinal tract including the esophagus, pylorus, duodenum, and colon.

Order Number	Reference Part Number	Catheter Fr	Catheter Length cm	Balloon Diameter mm	Balloon Fr	Balloon Length cm	Pressure atm	Pressure psi	Minimum Accessory Channel Diameter mm
G51799	HBD-W-8-9-10	8	240	8-9-10	24-27-30	5.5	4-6-8	60-90-120	2.8
G51795	HBD-W-10-11-12	8	240	10-11-12	30-33-36	5.5	2-4-6	30-60-90	2.8
G51796	HBD-W-12-13.5-15	8	240	12-13.5-15	36-40.5-45	5.5	2-4-6	30-60-90	2.8
G51797	HBD-W-15-16.5-18	8	240	15-16.5-18	45-49.5-54	5.5	2-4-6	30-60-90	2.8
G51798	HBD-W-18-19-20	8	240	18-19-20	54-57-60	5.5	2-4-6	30-60-90	2.8

Wire guide length is 300 cm.

Cook Sphere® Inflation Device

Used to inflate the balloon with fluid, to monitor the pressure in the balloon during the procedure, and to deflate the balloon after dilation.

Some products or part numbers may not be available in all markets. Contact your local Cook representative or Customer Support and Distribution for details regarding stock or nonstock status (expect longer lead time if the product is nonstock), pricing, and availability.

Customer Service

EU Website: cookmedical.eu EDI: cookmedical.eu/edi Distributors: +353 61239240, ssc.distributors@cookmedical.com Austria: +43 179567121, oe.orders@cookmedical.com Belgium: +32 27001702, be.orders@cookmedical.com Denmark: +45 38487607, da.orders@cookmedical.com Finland: +358 972519996, fi.orders@cookmedical.com France: +33 171230269, fr.orders@cookmedical.com Germany: +49 6950072804, de.orders@cookmedical.com Hungary: +36 17779199, hu.orders@cookmedical.com Iceland: +354 800 7615, is.orders@cookmedical.com Ireland: +353 61239252, ie.orders@cookmedical.com Italy: +39 0269682853, it.orders@cookmedical.com Netherlands: +31 202013367, nl.orders@cookmedical.com Norway: +47 23162968, no.orders@cookmedical.com Spain: +34 912702691, es.orders@cookmedical.com Sweden: +46 858769468, se.orders@cookmedical.com Switzerland - French: +41 448009609, fr.orders@cookmedical.com Switzerland - Italian: +41 448009609, it.orders@cookmedical.com Switzerland - German: +41 448009609, de.orders@cookmedical.com United Kingdom: +44 2073654183, uk.orders@cookmedical.com



USA Website: cookmedical.com EDI: cookmedical.com/edi.do Americas: Phone: +1 812.339.2235, 800.457.4500, Fax: 800.554.8335 E-mail: orders@cookmedical.com Australia: Phone: +61 734346000, 1800777222, Fax: +61 734346001, 1800077283 E-mail: cau.custserv@cookmedical.com

AI-ESC-IR-OHNS-PI-RH-SUR-A4